arXiv:math/0601288v1 [math.FA] 12 Jan 2006

RIESZ BASIS OF EXPONENTIALS FOR A UNION OF CUBES IN $\mathbb{R}^d$

JORDI MARZO

Abstract. We extend to several dimensions the result of K. Seip and Y.I. Lyubarskii that proves the existence of Riesz basis of exponentials for a finite union of intervals with equals lengths.

1. Introduction

Let $E \subset \mathbb{R}^d$ be a bounded set with no empty interior. We consider the Paley-Wiener space of L^2 functions band-limited to E

$$PW_E = \{F \in L^2(\mathbb{R}^d) : \sup \mathcal{F}F \subset E\},$$

endowed with the $L^2(\mathbb{R}^d)$ norm. Here we denote as

$$\mathcal{F}F(x) = \frac{1}{(2\pi)^{d/2}} \int_{\mathbb{R}^d} F(t) e^{-ix\cdot t} dt,$$

the Fourier transform of F.

By the Paley-Wiener theorem PW_E is a reproducing kernel Hilbert space of entire functions with kernel

$$K(z,w) = \frac{1}{(2\pi)^{d/2}} \int_E e^{it\cdot(z-\overline{w})} dt.$$

Then, for any $F \in PW_E$,

$$F(w) = \frac{1}{(2\pi)^{d/2}} \int_{\mathbb{R}^d} F(x) K(w,x) dx, \quad w \in \mathbb{C}^d.$$

Date: December 15, 2005.

2000 *Mathematics Subject Classification.* Primary 94A20; Secondary 30E05, 46E22, 30H05.

Key words and phrases. Sampling, Interpolation, Paley-Wiener spaces, Multiband functions, Riesz basis, multidimensional sampling, nonuniform sampling, Landau densities.

The author is supported by the DGICYT grant: BES-2003-2618, MTM2005-08984-C02-02 .

We say that a sequence $\Lambda = \{\lambda_n\} \subset \mathbb{R}^d$ is *sampling* for PW_E if there exist constants $A, B > 0$ such that

$$A \int_{\mathbb{R}^d} |F(x)|^2 dx \leq \sum_n |F(\lambda_n)|^2 \leq B \int_{\mathbb{R}^d} |F(x)|^2 dx, \tag{1}$$

for all $F \in PW_E$.

We say that a sequence $\Lambda = \{\lambda_n\} \subset \mathbb{R}^d$ is *interpolating* for PW_E if for each $\{a_n\} \in \ell^2$ there exists a function $F \in PW_E$ such that

$$F(\lambda_n) = a_n, \quad n \in \mathbb{Z}. \tag{2}$$

If Λ is both sampling and interpolating we say that it is a *complete interpolating sequence.*

Sampling and interpolation sequences in Paley-Wiener spaces provide a mathematical model of stable recovery and data transmission in signal theory regardless of the computation method. For the meaning of this concepts we refer to [Lan67b]. In 1967 H. J. Landau found necessary conditions for the existence of such a sequences in terms of some densities, as a consequence a complete interpolating sequence must have density equal to the Landau-Nyquist rate, see Theorem 2.1.

The aim of this paper is to find a complete interpolating sequence for PW_E, where E is a finite union of mutually disjoint half-open cubes

$$E = \bigcup_{j=1}^{p} Q_j \subset \mathbb{R}^d, \text{ where } Q_j = \prod_{s=1}^{d} [\alpha_j^s, \alpha_j^s + \beta), \ \ \beta > 0. \tag{3}$$

Our result allows us to construct, for any bounded E with no empty interior, sampling or interpolating sequences with density arbitrarily close to the critical one, see Corollary 3.4. This implies that Landau's results [Lan67] about densities can't be improved.

Moreover, we find a recovery process in PW_E whose stability is assured by the sampling property, see Theorem 3.3. This recovery process can be considered as a shifted version of the Shannon-Whittacker-Kotelnikov theorem.

The method used is a generalization to $\mathbb{R}^d$ with $d > 1$ to that of [LS97] for $\mathbb{R}$. It is worth noting that the reconstruction formula (14), which comes from A. Kohlenberg's work [Koh35], has been used by A. Faridani to produce sampling schemes for computerized tomography

[Far90]. Our result provides the stability that their result lacks. In fact, Faridani proved a more general version of the reconstruction formula (14) that holds for smooth functions with spectrum in any compact set of $\mathbb{R}^d$, and there is a version even for the group context [BF02]. But in such a general context there is no hope of stability.

Finally, we observe that we can define for $1 < q < \infty$ the Paley-Wiener space PW_E^q of entire functions given by

$$F(z) = \frac{1}{(2\pi)^{d/2}} \int_E f(t) e^{it\cdot z} dt, \quad z \in \mathbb{C}^d$$

with $f \in L^{q'}(E)$, for $q' = q/(q-1)$. With the natural modifications the main result (Theorem 3.3) holds also in PW_E^q for $1 < q < \infty$ because the projection of $L^q(\mathbb{R}^d)$ into $PW^q(E)$ is bounded and we have also a Plancherel-Pólya type inequality [Nik75, 3.2.2.].

2. Preliminaries

In this section we review some results and definitions that will be used later on.

Let $v = \{v_n\}$ be a sequence in a separable Hilbert space H. We say that v is a *frame* for H if there exist constants $A, B > 0$ such that

$$A||u||^2 \leq \sum_n |\langle u, v_n\rangle|^2 \leq B||u||^2, \quad u \in H.$$

We say that $v = \{v_n\}$ is a *Riesz sequence* or, equivalently, that it is a Riesz basis in its closed linear span, if there exist constants $A, B > 0$ such that for every finite scalar sequence $\{c_n\}$ one has

$$A \sum |c_n|^2 \leq \left\| \sum c_n v_n \right\|^2 \leq B \sum |c_n|^2.$$

Now we can translate our problem concerning sequences into the basis language. A sequence $\Lambda = \{\lambda_n\} \subset \mathbb{R}^d$ is interpolating for PW_E if and only if the sequence $\{K(\cdot, \lambda_n)/\sqrt{K(\lambda_n, \lambda_n)}\}$ is a Riesz sequence for PW_E, and this is also equivalent to the assumption that

$$\mathcal{E}(\Lambda) = \{e^{i\lambda_m \cdot x} : \lambda_m \in \Lambda\}$$

is a Riesz sequence for $L^2(E)$. Similarly, $\Lambda = \{\lambda_n\} \subset \mathbb{R}^d$ is sampling if and only if $\{K(\cdot, \lambda_n)/\sqrt{K(\lambda_n, \lambda_n)}\}$ is a frame for PW_E, and if and only if

$$\mathcal{E}(\Lambda) = \{e^{i\lambda_m \cdot x} : \lambda_m \in \Lambda\}$$

is a frame for $L^2(E)$. Then Λ is both sampling and interpolating if and only if $\mathcal{E}(\Lambda)$ is a Riesz basis for $L^2(E)$, i.e. $\mathcal{E}(\Lambda)$ is the image of an orthonormal basis by a bounded invertible operator, see [LS97].

We say that a sequence $\Lambda = \{\lambda_n\} \subset \mathbb{R}^d$ is *uniformly discrete* if

$$\inf_{j \neq i} |\lambda_j - \lambda_k| > 0.$$

This infimum is called the separation constant of Λ.

For $h > 0$ and $x = (x_1, \dots, x_d)$ denote by $Q_h(x)$ the cube:

$$Q_h(x) = \prod_{s=1}^{d} [x_s - h/2, x_s + h/2)\,.$$

Then $\{Q_h(hm)\}_{m \in \mathbb{Z}^d}$ is a disjoint cover of $\mathbb{R}^d$ for any $h > 0$.

Let $\Lambda = \{\lambda_n\} \subset \mathbb{R}^d$ be a sequence. Following Beurling we define the upper uniform density $D^+(\Lambda)$ and lower uniform density $D^-(\Lambda)$ of Λ as follows:

$$D^+(\Lambda) = \limsup_{h \to \infty} \frac{n^+(h)}{h^n} \quad \text{and} \quad D^-(\Lambda) = \liminf_{h \to \infty} \frac{n^-(h)}{h^n},$$

where

$$n^+(h) = \sup_{x \in \mathbb{R}^d} \#(\Lambda \cap Q_h(x)) \quad \text{and} \quad n^-(h) = \inf_{x \in \mathbb{R}^d} \#(\Lambda \cap Q_h(x)).$$

If this densities coincide we say that Λ has uniform density and we write $D(\Lambda) = D^+(\Lambda) = D^-(\Lambda)$.

The following result, which provides necessary conditions for sampling and interpolation, is due to H. J. Landau [Lan67].

Theorem 2.1. *If Λ is an interpolating sequence for PW_E then Λ is uniformly discrete and*

$$D^+(\Lambda) \leq \frac{|E|}{(2\pi)^d}.$$

If Λ is a sampling sequence for PW_E there exists $\Lambda' \subset \Lambda$ uniformly discrete and sampling subsequence such that

$$D^-(\Lambda') \geq \frac{|E|}{(2\pi)^d}.$$

Then Riesz basis can occur only when $D(\Lambda) = |E|/(2\pi)^d$. This particular value is called the Landau-Nyquist rate.

It is quite easy to show that no characterization can be obtained for sampling and interpolation in several dimensions, nor in the one-dimensional multi-band case, using only these densities.

Unlike the case of a single interval, where there is a complete characterization for the complete interpolating sequences [Pa79], little is known for the same problem with several intervals or in several dimensions. Let us comment some of the known results. The reasonable question in this more general setting is to find some complete interpolation sequence rather than to obtain a complete characterization. For a finite union of commensurable intervals, for two general intervals or for particular configurations with more than two intervals, it is known that such sequences exist [LS97]. In $\mathbb{R}^2$ there are also complete interpolating sequences for PW_E if E is a convex polygon symmetric with respect to the origin [LR00]. In $\mathbb{R}^d$ for $d > 2$ the only examples, as far as we know, concern the spaces PW_E with parallelogram domains E. For more on this problem see [Sei04].

Our result provides the first example of non-convex and non-symmetric set with Riesz basis of exponentials in several dimensions. This result contrast with the more rigid analogue problem of orthonormal basis of exponentials for $L^2(E)$, where it is known that symmetry is necessary [Kol00].

Our result allows us to construct, for any $E \subset \mathbb{R}^d$ bounded with no empty interior, sampling or interpolating sequences for PW_E with density arbitrarily close to the Landau-Nyquist rate, see Corollary 3.4. Unfortunately this method doesn't seem to be useful to find complete interpolating sequences for PW_E with E arbitrary.

Next we will show that the separation property of a sequence is equivalent to the right inequality in (1). This is a well known result but we include its proof for the sake of completeness [You80, pp 82-83].

Lemma 2.2. *Let $E \subset \mathbb{R}^d$ be a bounded set with no empty interior and $\Lambda = \{\lambda_n\} \subset \mathbb{R}^d$ be a sequence. Λ is a finite union of uniformly discrete sequences if and only if there exists a constant $B > 0$ such* *that*

$$\sum_n |F(\lambda_n)|^2 \leq B \int_{\mathbb{R}^d} |F(x)|^2 dx, \tag{4}$$

for all $F \in PW_E$. Moreover, the constant $B > 0$ depends only on the diameter of E and on the separation constant of Λ.

Proof. Suppose, to reach a contradiction, that we have (4) but Λ is not a finite union of uniformly discrete sequences. We can suppose that $0 \in \operatorname{int} E$ and then there exists an $\varepsilon > 0$ such that $Q_\varepsilon(0) \subset E$ and

$$\sup_{m \in \mathbb{Z}^d} \#(\Lambda \cap Q_h(hm)) = \infty,$$

where $h = \frac{\pi}{2\varepsilon}$. Choose a sequence $\{m_j\}_{j \in \mathbb{Z}} \subset \mathbb{Z}^d$ such that

$$\#(\Lambda \cap Q_h(hm_j)) \to \infty, \ \ j \to \infty.$$

The functions $f_j(x) = e^{-ihm_j \cdot x} \chi_{Q_\varepsilon(0)}(x)$ have support on E and

$$\begin{aligned} F_j(z) &= \mathcal{F}^{-1} f_j(z) = \frac{1}{(2\pi)^{d/2}} \int_{\mathbb{R}^d} f_j(x) e^{iz \cdot x} dx \\ &= \frac{1}{(2\pi)^{d/2}} \prod_{s=1}^d \frac{\sin \varepsilon (z^s - hm_j^s)}{z^s - hm_j^s}, \end{aligned}$$

are in PW_E. Now

$$\inf_{j \in \mathbb{Z}} \inf_{z \in Q_h(hm_j)} |F_j(z)| > 0,$$

because

$$\begin{aligned} \inf_{z \in Q_h(hm_j)} |F_j(z)| &= \frac{1}{(2\pi)^{d/2}} \inf_{z \in Q_h(0)} \left| \prod_{s=1}^d \frac{\sin \varepsilon z^s}{z^s} \right| \\ &\geq \left(\frac{\varepsilon}{\sqrt{2\pi}} \inf_{|t|<1/2} \left| \frac{\sin \pi t}{\pi t} \right| \right)^d, \end{aligned}$$

and therefore

$$||F_j|\Lambda||_{\ell^2} \to \infty, \ \ j \to \infty,$$

but $||F_j|| = |Q_\varepsilon(0)|^{1/2}$.

To prove the converse, we can suppose that Λ is uniformly discrete, i.e. there exists $\tau > 0$ such that

$$|\lambda_j - \lambda_k| \geq \tau > 0, \quad j \neq k.$$

Take an interval that contains E

$$E \subset \prod_{s=1}^{d} [-a^s, a^s].$$

Then $F \in PW_E$ is of exponential type $\nu = (\nu^1, \dots, \nu^d)$, where $0 < \nu^s \leq a^s$, $s = 1, \dots, p$. From the mean-value property one has, for all $\rho_s > 0$,

$$F(x) = \frac{1}{(2\pi)^d} \int_{[0,2\pi]^d} F(x^1 + \rho_1 e^{i\theta_1}, \dots, x^d + \rho_d e^{i\theta_d}) d\theta, \quad x \in \mathbb{R}^d.$$

Integrating on $[0, \delta]^d$, for some $\delta > 0$,

$$F(x) \left(\frac{\delta^2}{2} \right)^d = \frac{1}{(2\pi)^d} \int_{\prod_{s=1}^{d} B(x^s, \delta)} F(z) dm(z),$$

where $B(x^s, \delta) = \{z \in \mathbb{C} : |x^s - z| < \delta\}$. Then by Cauchy-Schwarz inequality

$$|F(x)|^2 \leq B \int_{[-\delta,\delta]^d} \left(\int_{\prod_{s=1}^{d} Q_{2\delta}(x^s)} |F(\xi + i\eta)|^2 d\xi \right) d\eta,$$

where $B > 0$ depends on δ and d.

For all F entire function of exponential type $(\nu^1, \dots \nu^d)$ such that $F \in L^2(\mathbb{R}^d)$, we have the following inequality of Plancherel-Pólya type

$$\|F(\cdot + i\eta)\|_{L^2(\mathbb{R}^d)} \leq e^{\sum_{s=1}^{d} \nu_s |\eta^s|} \|F\|_{L^2(\mathbb{R}^d)}, \tag{5}$$

see [Nik75, 3.2.2.].

Finally, if we take $\delta = \frac{\tau}{2}$ the cubes $Q_{2\delta}(\lambda_j)$ are pairwise disjoint. This together with (5), gives

$$\begin{aligned}\sum_j |F(\lambda_j)|^2 &\leq B\int_{[-\delta,\delta]^d}\sum_j \int_{\prod_{s=1}^d Q_{2\delta}(\lambda_j)}|F(\xi+i\eta)|^2 d\xi d\eta \\ &\leq B\int_{[-\delta,\delta]^d}\left(\int_{\mathbb{R}^d}|F(\xi+i\eta)|^2 d\xi\right)d\eta \\ &\leq B\|F\|^2_{L^2(\mathbb{R}^d)}\int_{[-\delta,\delta]^d} e^{\sum_{s=1}^d \nu_s|\eta^s|}d\eta \\ &= B\|F\|^2_{L^2(\mathbb{R}^d)}.\end{aligned}$$

■

Following Kohlenberg, [Koh35], we define the *sampling procedure* of order p for a function $F \in PW_E$ such that $\mathcal{F}F \in \mathcal{C}_c^\infty(E)$, as follows

$$G(t) = \sum_{l=1}^p \sum_{n\in\mathbb{Z}^d} F(Wn+k_l)S_l(t-Wn-k_l). \tag{6}$$

Here W is a real non-singular $d\times d$ matrix, $k_l \in \mathbb{R}^d$, and $S_l \in PW_E$. Since $\{Wn+k_l\}_{n\in\mathbb{Z}^d}$ is uniformly discrete (when the k_l are different) and F tends to zero faster than any polynomial this series converges. We want to determine W, k_l and S_l in order to get $G = F$ for all $F \in PW_E$.

First we compute the Fourier transform of this sampling procedure.

Proposition 2.3. *Let $F, S_l \in PW_E$, $f = \mathcal{F}F \in \mathcal{C}_c^\infty(E)$, $s_l = \mathcal{F}S_l$ and G given by (6). Then*

(7)
$$\mathcal{F}G(\omega) = \frac{(2\pi)^{d/2}}{|\det W|}\sum_{l=1}^p s_l(\omega)\sum_{n\in\mathbb{Z}^d} f(\omega+2\pi(W^{-1})^t n)e^{2\pi i (W^{-1})^t m\cdot k_l}.$$

Proof. This result follows easily from Poisson's Formula and it can be found in [Far90]. Consider the Fourier transform of G

$$\mathcal{F}G(\omega) = \sum_{l=1}^p s_l(\omega)\sum_{n\in\mathbb{Z}^d} F(Wn+k_l)e^{-i\omega\cdot(Wn+k_l)}.$$

Poisson's formula gives

$$(2\pi)^{d/2}|\det W|\sum_{n\in\mathbb{Z}^d} F_{k_l}(Wn)e^{-i\omega\cdot Wn}=\sum_{n\in\mathbb{Z}^d}\mathcal{F}F_{k_l}(\omega-2\pi(W^{-1})^t n)$$

where $F_{k_l}=F(\cdot+k_l)$. ■

3. Construction and main result

Let E be a finite union of half-open cubes

$$E=\bigcup_{j=1}^{p} Q_j\subset\mathbb{R}^d,\ \text{ where }\ Q_j=\prod_{s=1}^{d}[\alpha_j^s,\alpha_j^s+\beta),\ \ \beta>0.$$

We want to determine the functions and shifts $S_l(\cdot-k_l)$, $l=1,\dots,p$ in order that the reconstruction formula

$$F(t)=\sum_{l=1}^{p}\sum_{n\in\mathbb{Z}^d}F(Wn+k_l)S_l(t-Wn-k_l)$$

holds for all $F\in PW_E$ such that $f=\mathcal{F}F\in\mathcal{C}_c^\infty(E)$. We know, by Proposition 2.3, that for such functions this is equivalent to

$$(8)\quad f(\omega)=\frac{(2\pi)^{d/2}}{|\det W|}\sum_{l=1}^{p}s_l(\omega)\sum_{n\in\mathbb{Z}^d}f(\omega+2\pi(W^{-1})^t n)e^{2\pi i(W^{-1})^t n\cdot k_l}.$$

From now on we take $W=\frac{2\pi}{\beta}Id$ and, as $f=0$ off of E we define $s_l(\omega)=0$ if $\omega\notin E$. To determine s_l on E we split E in the parts which are overlapped by one or more of the sets

$$E+2\pi(W^{-1})^t n,\ \ n\in\mathbb{Z}^d.$$

Let us see how this partition works.

Consider two cubes Q_j and Q_k. There exists a unique $n_{jk}\in\mathbb{Z}^d$ such that

$$\gamma_{jk}=\alpha_k-\beta n_{jk}\in Q_j,$$

where $\alpha_k=(\alpha_k^1,\dots,\alpha_k^d)$ is the lower left point of Q_k. Observe that $n_{jj}=0$. Fixed $j\in\{1,\dots,p\}$ split Q_j into the half-open intervals formed by intersection of Q_j with the 2^d quadrants around γ_{jk}, and do the same for all of $k=1,\dots,p$. Thus, we split each cube Q_j in this way in at most p^d intervals (this can be easily proved by induction, projecting a face of the cube into $\mathbb{R}^{d-1}$) which we denote Q_j^s $s=$



FIGURE 1.

$1,\dots,p^d$, the subscript indicates the cube in which the rectangle is contained. From now on we assume that we have p^d intervals Q_j^s on each cube Q_j. The minor changes needed if that is not the case are left to the reader. We have for each $j=1,\dots,p$

$$\bigcup_{s=1}^{p^d} Q_j^s = Q_j, \quad Q_j^s \cap Q_j^\ell = \emptyset.$$

Although $\gamma_{jk}+\beta n_{jk}=\alpha_k \in Q_k$ and $\gamma_{jk}\in Q_j$, it can happen that for some $Q_j^s \subset Q_j$

$$(Q_j^s+\beta n_{jk})\cap Q_k=\emptyset,$$

see figure 1. To solve this difficulty we choose a correction

$$C_{js}(k)\in\{-1,0,1\}^d$$

in such a way that

$$Q_j^s+\beta(n_{jk}+C_{js}(k)) = Q_k^{l_s},$$

for some $l_s\in\{1,\dots,p^d\}$.

This shows that the intervals that form the cubes are always the same, but put in different orders, i.e. for any $Q_1^s\subset Q_1$ we have a unique rectangle in Q_j that is translated by $\beta\mathbb{Z}^d$ in Q_1^s. We call this rectangle Q_j^s. Then the intervals on each Q_j are named depending on their translate on Q_1. Using this convention we have for $s = 1, \ldots, p^d$ $k = 1, \ldots, p$

$$Q_1^s + \beta(n_{1k} + C_{1s}(k)) = Q_k^s,$$

and

$$n_{kj} + C_{ks}(j) = n_{1j} - n_{1k} + C_{1s}(j) - C_{1s}(k). \tag{9}$$

This is so because the translation by $\beta\mathbb{Z}^d$ that moves Q_j^s to their counterpart in Q_k is unique.

Let

$$C_{js} = \{n_{jk} + C_{js}(k)\}_{k=1,\ldots p}$$

and notice that $0 = n_{jj} + C_{js}(j) \in C_{js}$ and as the cubes are disjoint the elements in C_{js} are all different.

Now we have, for $\omega \in Q_j^s$

$$\omega + \beta n \in E \quad \text{iff} \quad n \in C_{js},$$

then for $\omega \in Q_j^s$ equation (8) becomes

$$\begin{aligned} f(\omega) &= \left(\frac{\beta}{\sqrt{2\pi}}\right)^d \sum_{l=1}^{p} s_l(\omega) \sum_{n \in C_{js}} f(\omega + \beta n) e^{i\beta n \cdot k_l} \\ &= \left(\frac{\beta}{\sqrt{2\pi}}\right)^d \sum_{n \in C_{js}} f(\omega + \beta n) \sum_{l=1}^{p} s_l(\omega) e^{i\beta n \cdot k_l}. \end{aligned}$$

If we find for all j and s a solution $x_l = x_l(j,s)$ $l = 1, \ldots, p$ of the linear system

$$\begin{cases} \sum_{l=1}^{p} x_l e^{i\beta m \cdot k_l} = 0, & \text{if } m \in C_{js} \setminus \{0\} \\ \sum_{l=1}^{p} x_l = \left(\frac{\beta}{\sqrt{2\pi}}\right)^{-d}, & \text{otherwise} \end{cases} \tag{10}$$

we have finished, because defining $s_l(\omega) = x_l(j,s)$ for $\omega \in Q_j^s$ and

$$s_l(\omega) = \sum_{j=1}^{p} \sum_{s=1}^{p^d} x_l(j,s) \chi_{Q_j^s}(\omega)$$

we have (8).

We claim that there exist $k_l \in \mathbb{R}^d$ for $l = 1, \ldots, p$ such that (10) has a solution for all $j = 1, \ldots, p$ and $s = 1, \ldots, p^d$.

Indeed, letting $z_{ls} = e^{i\beta k_l^s}$ for $l = 1, \dots, p$, $s = 1, \dots, d$ and $z_l = (z_{l1}, \dots, z_{ld})$ the linear system in (10) has solution if and only if there exists $z_l \in \mathbb{T}^d$, $l = 1, \dots, p$ such that

$$\mathcal{D}(z_{11}, \dots, z_{pd}) = \prod_{j=1}^{p} \prod_{s=1}^{p^d} \det(\{z_l^{n_{jk}+C_{js}(k)}\}_{k,l=1,\dots,p}) \neq 0. \tag{11}$$

Indeed, letting

$$m_{js} = (\min_{k=1,\dots,p} n_{jk}^1 + C_{js}^1(k), \dots, \min_{k=1,\dots,p} n_{jk}^d + C_{js}^d(k)),$$

then

$$\det(\{z_l^{n_{jk}+C_{js}(k)}\}_{k,l=1,\dots,p}) = \frac{pol_{js}(z_{11}, \dots, z_{pd})}{z_1^{m_{js}} \cdots z_p^{m_{js}}}$$

where pol_{js} is a non-zero polynomial in $\mathbb{C}^{pd}$. Now the first column expansion of the determinant gives us

$$\sum_{k=1}^{p} (-1)^k z_1^{n_{jk}+C_{js}(k)} M_{1k}(z_2, \dots, z_p).$$

Since the elements in C_{js} are different, an induction reasoning provides the conclusion.

Moreover, the zero set of $\mathcal{D}$ has zero Lebesgue measure in the polydisc $\mathbb{T}^{pd}$, because $\mathcal{D}$ is an entire function [Ron74, p. 218]. Then for almost all $(k_1, \dots, k_p) \in [0, \frac{2\pi}{\beta})^{pd}$, and all j and s the linear system in (10) has a solution.

So far we have shown that for $F \in PW_E$ such that $\mathcal{F}F \in \mathcal{C}_c^\infty(E)$, there exist $k_l \in \mathbb{R}^d$ with

$$\mathcal{D}(e^{i\beta k_1^1}, \dots, e^{i\beta k_p^d}) \neq 0, \tag{12}$$

so that taking

$$S_l(t) = \mathcal{F}^{-1} s_l(t) = \sum_{j=1}^{p} \sum_{s=1}^{p^d} x_l(j,s) \mathcal{F}^{-1} \chi_{Q_j^s}(t), \quad l = 1, \dots, p, \tag{13}$$

we have

$$F(t) = \sum_{l=1}^{p} \sum_{n \in \mathbb{Z}^d} F(Wn + k_l) S_l(t - Wn - k_l). \tag{14}$$

The following results allows us to extend this formula to all $F \in PW_E$.

Lemma 3.1. *Take $k_l \in \mathbb{R}^d$ such that (12) holds and functions S_l as in (13). Then, if $\alpha = \cup_{l=1}^p \alpha^l$, $\alpha^l \in \ell^2$, the series*

$$F[\alpha](t) = \sum_{l=1}^{p} \sum_{n \in \mathbb{Z}^d} \alpha_n^l S_l(t - Wn - k_l), \tag{15}$$

converge in $L^2(\mathbb{R}^d)$. Moreover $F[\alpha]$ is an injective operator: there exists $B > 0$ independent of α such that

$$\|F[\alpha]\|_{L^2(\mathbb{R}^d)} \leq B\|\alpha\|_{\ell^2}. \tag{16}$$

Corollary 3.2. *Taking functions S_l as in (13), and $k_l \in \mathbb{R}^d$ such that (12) holds, then*

$$S_l(Wn + k_s - k_l) = \delta_{n,0}\delta_{l,s}, \quad l, s = 1, \ldots, p.$$

The proofs are given in the next section. We can now prove our main result.

Theorem 3.3. *Let $E \subset \mathbb{R}^d$ be a finite union of cubes as in (3). If $W = \frac{2\pi}{\beta} Id$ and $k_l \in \mathbb{R}^p$ are such that (12) holds then*

$$\Lambda(k_1, \ldots, k_p) = \cup_{l=1}^p \{Wn + k_l\}_{n \in \mathbb{Z}^d}$$

is a complete interpolating sequence for PW_E, i.e. there exists constants $A, B > 0$ such that

$$A\|F\|^2 \leq \sum_{l=1}^{p} \sum_{n \in \mathbb{Z}^d} |F(Wn + k_l)|^2 \leq B\|F\|^2, \quad F \in PW_E. \tag{17}$$

Given any $\alpha = \cup_{l=1}^p \alpha^l$ with $\alpha^l \in \ell^2$ there exists a unique $F \in PW_E$ such that

$$F(Wn + k_l) = \alpha_n^l, \quad n \in \mathbb{Z}^d, l = 1, \ldots, p. \tag{18}$$

Moreover, taking functions S_l as in (13), the reconstruction formula (14) holds for all $F \in PW_E$, with convergence in L^2-norm and uniform convergence on products of horizontal strips.

Proof. By the preceding construction we know that (14) holds for all $F \in PW_E$ such that $\mathcal{F}F \in \mathcal{C}_c^\infty(E)$. Now by Lemma 3.1 the operator on PW_E defined by

$$F \longmapsto \sum_{l=1}^{p} \sum_{n \in \mathbb{Z}^d} F(Wn + k_l) S_l(t - Wn - k_l),$$

is bounded and the sampling formula (14) holds for PW_E.

Since $\Lambda(k_1, \ldots, k_p)$ is uniformly discrete we have the right inequality in (17). In order to prove the left inequality, notice that by Lemma 3.1 we have that $\sum \alpha_n^l S_l(\cdot - Wn - k_l)$ converges for all $\alpha^l \in \ell^2$, therefore there exist a constant $B > 0$ such that for $F \in PW_E$

$$\sum_{l,n} |\langle F, S_l(\cdot - Wn - k_l) \rangle|^2 \leq B \|F\|^2,$$

see [Chr03, Lemma 3.2.1.]. Then for $F \in PW_E$ we have

$$\begin{aligned} \|F\|^4 &\leq \left(\sum_{l,n} |F(Wn + k_l)|^2 \right) \left(\sum_{l,n} |\langle F, S_l(\cdot - Wn - k_l) \rangle|^2 \right) \\ &\leq B \|F\|^2 \left(\sum_{l,n} |F(Wn + k_l)|^2 \right). \end{aligned}$$

Defining for $\alpha = \cup_{l=1}^{p} \alpha^l$, $\alpha^l \in \ell^2$, $F[\alpha] \in PW_E$ as in (15), and using Corollary 3.2 we get the interpolation property (18)

$$\begin{aligned} F[\alpha](Wm + k_s) &= \sum_{l=1}^{p} \sum_{n \in \mathbb{Z}^d} \alpha_n^l S_l(W(m-n) + k_s - k_l) \\ &= \sum_{l=1}^{p} \sum_{n \in \mathbb{Z}^d} \alpha_n^l \delta_{m,n} \delta_{l,s} = \alpha_m^s. \end{aligned}$$

Finally, the uniform convergence on products of horizontal strips

$$\{z \in \mathbb{C}^d : |\Im z^j| \leq C_j, j = 1, \ldots, p\},$$

follows from the Phragmén-Lindelöf inequality, see [You80, p 84]. ■

Corollary 3.4. *Let $E \subset \mathbb{R}^d$ be a bounded set with no empty interior. For all $\varepsilon > 0$ there exist uniformly separated sequences $\Lambda^\varepsilon, \Lambda_\varepsilon \subset \mathbb{R}^d$ such that Λ_ε is sampling for PW_E, Λ^ε is interpolating for PW_E and*

$$0 \leq (2\pi)^d D(\Lambda_\varepsilon) - |E| < \varepsilon, \quad 0 \leq |E| - (2\pi)^d D(\Lambda^\varepsilon) < \varepsilon.$$

Proof. It is enough to choose E_ε, E^ε as in (3) such that $E^\varepsilon \subset E \subset E_\varepsilon$ and $|E_\varepsilon| - |E|, |E| - |E^\varepsilon| < \varepsilon$. Now we take a complete interpolating sequence Λ^ε for PW_{E^ε} (that obviously is also interpolating for PW_E) and Λ_ε a complete interpolating sequence for PW_{E_ε} (that is also sampling for PW_E) and apply Landau's Theorem 2.1. ■

4. PROOFS OF TECHNICAL RESULTS

In this section we will proof the technical results used in proving Theorem 2.3.

Proof. [Lemma 3.1] Consider, given $M \in \mathbb{N}$

$$F_M[\alpha](t) = \sum_{l=1}^{p} \sum_{|n| \leq M} \alpha_n^l S_l(t - Wn - k_l).$$

For all $f \in L^2(\mathbb{R}^d)$ we have

$$\begin{aligned}
\langle F_M[\alpha], f \rangle &= \sum_{l,j=1}^{p} \sum_{s=1}^{p^d} x_l(j,s) \sum_{|n| \leq M} \alpha_n^l \langle e^{-i\omega \cdot (\frac{2\pi}{\beta} n + k_l)}, \chi_{Q_j^s} \widehat{f} \rangle \\
&= \sum_{l,j=1}^{p} \sum_{s=1}^{p^d} x_l(j,s) \sum_{|n| \leq M} \alpha_n^l \mathcal{P}_{PW_{Q_j^s}}(f) \left(\frac{2\pi}{\beta} n + k_l \right),
\end{aligned}$$

where $\mathcal{P}_{PW_{Q_j^s}}(f) = \mathcal{F}^{-1}(\chi_{Q_j^s}\mathcal{F}f)$ is the orthogonal projection of $f \in L^2(\mathbb{R}^d)$ into $PW_{Q_j^s}$. Then, using Lemma 2.2 (d), we obtain

$$\begin{aligned}|\langle F_M[\alpha], f\rangle| &\leq \sum_{l,j=1}^{p}\sum_{s=1}^{p^d}|x_l(j,s)|\|\alpha^l\|_{\ell^2}\|\mathcal{P}_{PW_{Q_j^s}}(f)\|_{L^2}\\ &\leq B\|f\|_{L^2}\sum_{l=1}^{p}\|\alpha^l\|_{\ell^2}\sum_{j=1}^{p}\sum_{s=1}^{p^d}|x_l(j,s)|\|\mathcal{P}_{PW_{Q_j^s}}\|\\ &\leq B\|f\|_{L^2}\sum_{l=1}^{p}\|\alpha^l\|_{\ell^2},\end{aligned}$$

where the constant B does not depend on α.

Now we want to prove the injectivity part. By taking Fourier transform in (15) we obtain

$$0 = \sum_{j=1}^{p}\sum_{s=1}^{p^d}\chi_{Q_j^s}(\omega)\sum_{l=1}^{p}x_l(j,s)e^{-i\omega\cdot k_l}\left\{\sum_{n\in\mathbb{Z}^d}\alpha_n^l e^{-i\frac{2\pi}{\beta}\omega\cdot n}\right\}. \tag{19}$$

We call

$$f_l(\omega) = \sum_{n\in\mathbb{Z}^d}\alpha_n^l e^{-i\frac{2\pi}{\beta}\omega\cdot n}, \quad l = 1,\dots,p.$$

If we show that

$$f_l(\omega) = 0, \quad l = 1,\dots p,$$

we are done. The functions f_l are $\beta-$periodic on each variable, hence it is enough to proof $f_l|Q_1 = 0$. We will prove this on each Q_1^s separately. The key is property (9).

Let us take $\omega \in Q_1^s$. Then, as $f_l(\omega + \beta(n_{1k} + C_{1s}(k))) = f_l(\omega)$ and $\omega + \beta(n_{1k} + C_{1s}(k)) \in Q_{ks}$, substituting into (19) we obtain, for $k = 1,\dots,p$

$$\begin{aligned}0 &= \sum_{l=1}^{p}x_l(j,s)e^{-i(\omega+\beta(n_{1k}+C_{1s}(k)))\cdot k_l}f_l(\omega)\\ &= \sum_{l=1}^{p}x_l(j,s)e^{-i\beta(n_{1k}+C_{1s}(k))\cdot k_l}[e^{-i\omega\cdot k_l}f_l(\omega)].\end{aligned}$$

We have a linear system with unknowns $e^{-i\omega\cdot k_l} f_l(\omega)$ and coefficient's matrix

$$(x_l(k,s)e^{-i\beta(n_{1k}+C_{1s}(k))\cdot k_l})_{k,l} = (a_{kl})_{k,l}.$$

We will see that this system has only the trivial solution by showing that it has invertible coefficient's matrix. This point becomes quite easy with the notation we have adopted. Indeed, the linear system (10) that we have used to determine s_l in Q_1^s is given by

$$(e^{i\beta(n_{1j}+C_{1s}(j))\cdot k_l})_{l,j} = (b_{lj})_{l,j}.$$

Now by (9), the elements of the matrix product are

$$\begin{aligned}\sum_{l=1}^{p} a_{kl}b_{lj} &= \sum_{l=1}^{p} x_l(k,s)e^{i\beta(n_{1j}-n_{1k}+C_{1s}(j)-C_{1s}(k))\cdot k_l} \\ &= \sum_{l=1}^{p} x_l(k,s)e^{i\beta(n_{kj}+C_{ks}(j))\cdot k_l} = \delta_{kj}\left(\frac{\sqrt{2\pi}}{\beta}\right)^d\end{aligned}$$

which is the desired conclusion. ■

Proof. [Corollary 3.2] Let $s \in \{1,\dots,p\}$ be fixed. The sampling result applied to $S_s(\cdot - k_l) \in PW_E$ yields

$$S_s(t-k) = \sum_{l=1}^{p}\sum_{n\in\mathbb{Z}^d} S_s(Wn + k_l - k_s)S_l(t - Wn - k_l).$$

Now applying Lemma 3.1 to

$$\sum_{l=1}^{p}\sum_{n\in\mathbb{Z}^d}\left(\delta_{n,0}\delta_{l,s} - S_s(Wn+k_l)\right)S_l(t-Wn-k_l) = 0,$$

we obtain the result. ■

References

[Far90] A. Faridani, *An application of a multidimensional theorem to computed tomography*, Contemp. Math. 113, 65–80, 1990

[BF02] H. Behmard, A. Faridani, *Sampling of bandlimited functions on unions of shifted lattices*, J. Fourier Anal. Appl. , vol. 8, **1**, 2002

[Koh35] A. Kohlenberg, *Exact interpolation of Band-Limited Functions*, J.Appl.Phys., **24**, 12, 1432–1436, 1935

[Kol00] M. Kolountzakis, *Non-symmetric convex domains have no basis of exponentials*, Illinois J. Math., **44**, no. 3 , 542–550, 2000

[Lan67] H. J. Landau, *Necessary Density Conditions for Sampling and Interpolation of Certain Entire Functions*, Acta. Math., **117**, 37–52, 1967

[Lan67b] H. J. Landau, *Sampling, Data transmission, and the Nyquist rate*, Proc. IEEE, vol. 55, no. 10, 1701–1706, 1967

[LR00] Y. I. Lyubarskii, A. Rashkovskii, *Complete interpolating sequences for Fourier transforms supported by convex symmetric polygons*, Ark. Mat. **38**, 139–170, 2000

[LS97] Y. I. Lyubarskii, K. Seip, *Sampling and Intepolating Sequences for Multiband-Limited Functions end Exponential Bases on Disconnected Sets*, J. Fourier Anal. Appl., **3**, 5, 597-615, 1997

[Nik75] S. M. Nikol'skiĭ, *Approximation of Functions of Several Variables and Imbedding Theorems*, Springer-Verlag, Die Grundlehren der mathematische Wissenschaften in Einzeldarstellungen, 205, 1975

[Chr03] O. Christensen, An introduction to frames and Riesz basis, Birkhäuser, ANHA, 2003

[Pa79] B. S. Pavlov, *The basis property of a system of exponentials and the condition of Muckenhoupt*, Dokl. Akad. Nauk SSSR, **247**, no. 1, 37–40, 1979

[Ron74] L. I. Ronkin, *Introduction to the theory of entire functions of several variables*, Trans. Math. Monographs, AMS, Providence, RI, 1974

[Sei04] K. Seip, *Interpolation and sampling in spaces of analytic functions. University Lecture Series*, 33. American Mathematical Society, Providence, RI, 2004

[You80] R. Young, *An introduction to nonharmonic Fourier series*, Academic Press, New York, 1980 (revised first edition 2001)

Departament de matemàtica aplicada i anàlisi, Universitat de Barcelona, Gran via 585, 08071 Barcelona, Spain

E-mail address: `jmarzo@mat.ub.es`